

# ON THE DENOMINATOR OF THE POINCARÉ SERIES FOR MONOMIAL QUOTIENT RINGS

HARA CHARALAMBOUS

ABSTRACT. Let  $S = \mathbb{k}[x_1, \dots, x_n]$  be a polynomial ring over a field  $\mathbb{k}$  and  $I$  a monomial ideal of  $S$ . It is well known that the Poincaré series of  $\mathbb{k}$  over  $S/I$  is rational. We describe the coefficients of the denominator of the series and study the multigraded homotopy Lie algebra of  $S/I$ .

## CONTENTS

|                                                |   |
|------------------------------------------------|---|
| 1. Introduction                                | 1 |
| 2. The monomials of $Q_R(y, t)$                | 2 |
| 3. Multigraded deviations and acyclic closures | 4 |
| References                                     | 6 |

## 1. INTRODUCTION

Let  $S = \mathbb{k}[x_1, \dots, x_n]$  be a polynomial ring over a field  $\mathbb{k}$ ,  $I$  a monomial ideal of  $S$  and  $R = S/I$ . If  $J$  is a subset of the minimal monomial generating set  $I$  of  $I$ , we let  $m_J$  denote the least common multiple of the monomials in  $J$  or the corresponding monomials in  $\mathbb{k}[y_1, \dots, y_n]$  as appropriate. The multigraded Poincaré series of  $R$ ,  $P_R(y, t)$  is rational, [Ba82]:

$$P_R(y, t) = \sum_{i=0}^{\infty} \sum_{j \geq 0} y^j \dim_k(\mathrm{Tor}_i^R(k, k)_j) t^i = \frac{\prod_{i=1}^n (1 + ty_i)}{Q_R(y, t)}$$

where  $Q_R(y, t) \in Z[y_1, \dots, y_n][t]$ .

The few facts that are known in general about the multigraded expansion  $Q_R(y, t)$  also date back to [Ba82]: the degree of  $Q_R(y, t)$  in  $t$  is bounded above by the total degree of  $m_I$ , and the monomial coefficients of  $t$  divide  $m_I$ . In this paper we discuss the monomial coefficients of  $Q_R(y, t)$ .

In the first section of this paper we show that the monomial coefficients of  $t$  in  $Q_R(y, t)$  are least common multiples of the monomial generators of  $I$ . We discuss the Koszul homology of  $R$  when  $R$  is Golod. We also present a free resolution of  $R$  when  $I$  is generic. In the second section we discuss the multigraded acyclic closure of  $R$  as well as the multigraded deviations in terms of the multigraded homotopy Lie algebra of  $R$ , and show that the LCM lattice determines the Poincaré series of  $R$ . This is the multigraded version of Theorem 1 of [Av02].

Thanks are due to J. Backelin and the faculty of the Department of Mathematics in Stockholm, for useful discussions during my brief stay in 1996.

---

*Date:* February 1, 2008.

2. THE MONOMIALS OF  $Q_R(y, t)$ 

The argument used in the proof of the next proposition has the flavor of [Ba82] and uses Golod theory. It uses Lescot's result, [Le86], that the Poincaré series  $M$  over  $R$

$$P_R^M(y, t) = \sum_{i=0}^{\infty} \sum_{\mathbf{j} \geq 0} y^{\mathbf{j}} \dim_k(\mathrm{Tor}_i^R(M, k)_{\mathbf{j}}) t^i = \frac{f_R^M(y, t)}{Q_R(y, t)}$$

where  $f_r^M(y, t) \in Z[y_1, \dots, y_n][t]$ .

**Proposition 2.1.** *Let  $Q_R(y, t) = 1 + \sum(\sum c_{\mathbf{j}} y^{\mathbf{j}}) t^i$ . Then  $y^{\mathbf{j}}$  is equal to a least common multiple of a subset of the minimal monomial generating set of  $I$ .*

*Proof.* By a standard argument, [Ba82], it is enough to prove Proposition 2.1 when the ideal  $I$  is generated by squarefree monomials. In this case the  $y$  monomials of  $Q_R(y, t)$  are squarefree, [Ba82]. We do induction on the number of variables  $n$  that divide the monomials of the minimal generating set of  $I$ , the case  $n = 1$  being trivial. Consider a term of  $Q_R(y, t)$  whose coefficient in  $y$  does not involve  $y_i$ . We separate the generators of  $I$  that involve  $x_i$ . Let  $I = L_i + x_i J_i$  where  $L_i, J_i \subset \mathbb{k}[x_1, \dots, \hat{x}_i, \dots, x_{n-1}]$  and let  $A = \mathbb{k}[x_1, \dots, \hat{x}_i, \dots, x_{n-1}]/L_i$  and  $A' = \mathbb{k}[x_1, \dots, \hat{x}_i, \dots, x_{n-1}]/(L_i + J_i)$ . Therefore  $R \cong A[x_i]/x_i J_i$  and  $A[x_i] \rightarrow R$  is Golod, which implies that

$$\begin{aligned} P_R(y, t) &= \frac{P_{A[x_i]}}{(1 - t(P_{A[x_i]}^R - 1))} = \frac{(1 + ty_i)P_A}{(1 - t^2 P_{A[x_i]}^{x_i J_i})} = \frac{(1 + ty_i)P_A}{(1 - t^2 y_i P_A^{J_i})} = \\ &= \frac{(1 + ty_i)P_A}{(1 - ty_i(P_A^{A'} - 1))} = \frac{\prod(1 + ty_i)}{Q_A + ty_i Q_A - ty_i f_A^{A'}}. \end{aligned}$$

The  $y$  monomials that do not involve  $y_i$  are terms of  $Q_A$ , so by induction they have the desired form. Suppose now that a monomial of  $Q_R(y, t)$  involves all  $y_i$ : it is the product of all the variables  $y_i$  and is equal to the least common multiple of all the generators in  $I$ .  $\square$

In [ChRe95] we noticed that the terms of  $Q_R(y, t)$  when  $I$  is a complete or almost complete intersection come from the set  $\{(-1)^{l_{\mathbf{j}}} t^{(|\mathbf{j}| + l_{\mathbf{j}})} m_{\mathbf{j}}\}$ . Here  $\mathbf{j}$  ranges over the subsets of the minimal monomial generating set of  $I$  and  $l_{\mathbf{j}}$  is the number of connected components of the graph we get when we connect the elements of  $\mathbf{j}$  by an edge if they have a variable in common. In [ChRe95] we also confirmed that this is the case when the Taylor resolution [Ta60], [Ei97] of  $S/I$  over  $S$  is minimal.

In Remark 2.3 we confirm that the same holds for the monomial ideals  $I$  such that  $R/I$  is Golod. First we recall and expand a comment by Fröberg, [Fr79].

**Remark 2.2.** Let  $R = S/I$  be Golod and  $K_{\bullet}$  be the Koszul complex on the variables  $\bar{x}_i$  of  $R$ . Then  $H(K_{\bullet})$  is isomorphic to  $\mathbb{k}[X_{A_1}, \dots, X_{A_r}]/(X_{A_1}, \dots, X_{A_r})^2$  where  $A_i$  is a subset of the minimal monomial generating set of  $I$  and the degree of  $X_{A_i}$  equals the cardinality of  $A_i$ .

Indeed since  $K_{\bullet}$  admits trivial Massey operations the product of two elements of  $H_{\geq 1}(K_{\bullet})$  is zero, [GuLe69], [Av98]. Moreover  $H(K_{\bullet})$  is always of the form  $\mathbb{k}[X_1, \dots, X_r]/L$ :  $L$  is generated by quadratics, the  $X_i$ 's form a minimal multiplicative generating set for  $H(K_{\bullet})$ , and  $X_i X_j = (-1)^{|X_i||X_j|} X_i X_j$  where  $|X_i|$  is the degree of  $X_i$ . Let  $F_{\bullet}$  be the Taylor complex on the minimal monomial generators of  $I$  over  $S$ , [Ta60] and let  $E_{\bullet}$  be the minimal multigraded resolution of  $S/I$  over

$Q$ .  $E_\bullet$  is a direct summand of  $F_\bullet$ . It follows that the multidegrees of the generators of  $E_i$  are among the multidegrees of the generators of  $F_i$ . Therefore they are the multidegrees of least common multiples of subsets  $A_j$  of  $I$ . It follows that  $H(K_\bullet) = H(\mathbb{k} \otimes E_\bullet) = \mathbb{k}[X_{A_1}, \dots, X_{A_r}]/L$ . Since  $X_{A_j}X_{A_t} = 0$ ,  $X_{A_j}X_{A_t} \in L$ .

**Remark 2.3.** Thus monomial Golod rings are special Golod rings. Let  $R$  be a Golod ring. From our previous remark and using the fact that  $m_A \neq m_{A_1} \cdots m_{A_t}$  for any partition of  $A$  where  $X_A$  and  $X_{A_j}$  are among the generators of  $H(K_\bullet)$ , we see that  $A$  is connected and  $l_A = 1$ . Let  $H_i(K)_j$  denote the subspace of  $H_i(K)$  of multidegree  $j$ . Thus the terms of the denominator of the Poincaré series of  $R$ ,  $Q_R(y, t) = 1 - \sum_i \sum_j \dim_{\mathbb{k}} H_i(K)_j y^j t^{i+1}$  of the Golod ring  $R$  come from the set  $\{(-1)t^{(|J|+1)}m_J\}$  where  $J$  ranges over the subsets of the minimal monomial generating set of  $I$ .

Generic ideals and their minimal resolutions were first introduced in [BaPeSt98] and generalized in [MiStYa00]. Let  $I$  be a generic ideal. If the multidegrees of two minimal monomial generators of  $I$  are equal for some variable then there is a third monomial generator of  $I$  whose multidegree is strictly smaller than the multidegree of the least common multiple of the other two. The minimal resolution  $E_\bullet$  of  $S/I$  over  $S$  is determined by the Scarf complex of  $\Delta_I$ . The latter consists of the subsets  $J$  of the index set of  $I$ ,  $\Phi$ , such that  $m_J \neq m_L$  for any other subset  $L$  of  $\Phi$ . We note that  $E_\bullet$  has the structure of a DG algebra, [BaPeSt98]. The Eagon resolution of  $\mathbb{k}$  over  $R = S/I$  was discussed in [GuLe69]. Below we give the generators for the Eagon resolution when  $I$  is generic.

**Theorem 2.4.** *Let  $R = S/I$  where  $I$  is a generic ideal of  $S$ , let  $K_\bullet$  be the Koszul complex on the  $x_i$  over  $R$ ,  $X_0 = 0$ , and  $X_i$  be the free  $R$ -module with generators  $T_L$  where  $|L| = i$  and  $L$  is in  $\Delta_I$ , the Scarf complex of  $I$ . Then  $(Y_\bullet, d_\bullet)$  is a free resolution of  $\mathbb{k}$  over  $R$ , where*

$$Y_i = \sum_{j+t_1+\dots+t_l=i} K_j \otimes X_{t_1} \otimes \cdots \otimes X_{t_l},$$

and

$$d(T_J) = (m_J/x_J)e_J,$$

$$d(T_{I_1} \otimes \cdots \otimes T_{I_l}) = d(T_{I_1} \otimes \cdots \otimes T_{I_{l-1}}) \otimes T_{I_l} +$$

$$\sum_{i=1}^{l-1} (|I_i| + 1) (-1)^{i-1} T_{I_1} \otimes \cdots \otimes T_{I_{l-2}} \otimes \delta(I_{l-1}, I_l) (T_{I_{l-1}} \wedge T_{I_l}),$$

where  $\delta(I_{l-1}, I_l) = \frac{m_{I_{l-1}}m_{I_l}}{m_{I_{l-1} \cup I_l}}$  if  $I_{l-1} \cup I_l \in \Delta_I$  and 0 otherwise

$$d(e_L \otimes T_{I_1} \otimes \cdots \otimes T_{I_l})d(e_L) \otimes T_{I_1} \otimes \cdots \otimes T_{I_l} + (-1)^{|L|} e_L \otimes d(T_{I_1} \otimes \cdots \otimes T_{I_l}).$$

*Proof.* The theorem follows immediately from [Iy97], Theorem 1.2 and the preceding comments.  $\square$

As an immediate corollary we get the characterization of Golod rings when  $I$  is a generic ideal, which generalizes Proposition 2.4 of [Ga00].

**Proposition 2.5.** *Let  $I$  be a generic ideal.  $S/I$  is Golod if and only if  $m_{I_1}m_{I_2} \neq m_{I_1 \cup I_2}$  whenever  $I_1 \cup I_2 \in \Delta_I$ .*

## 3. MULTIGRADED DEVIATIONS AND ACYCLIC CLOSURES

In this section we introduce the multigraded acyclic closure  $R\langle X \rangle$  of  $\mathbb{k}$  over  $R$ . For the construction of the usual acyclic closure, we refer to Construction 6.3.1, [Av98]. To this end we choose  $x_j$ , the  $\Gamma$ -variables of  $X_n$  so that  $\text{cls}(\theta(x_j))$  has multidegree  $\mathbf{j}$  and  $\{\text{cls}(\theta(x_j))\}$  is a minimal multigraded generating set of  $H_{n-1}(R\langle X_{\leq n-1} \rangle)$ . We define the multidegree of  $x_j$  to be  $\mathbf{j}$  and we let  $X_{n,\mathbf{j}}$  consist of all  $x \in X_n$  whose multidegree is  $\mathbf{j}$ . We will show that the cardinality of  $X_{n,\mathbf{j}}$  appears as an exponent in a product decomposition of the multigraded Poincaré series.

First we state and prove the multigraded version of Remark 7.1.1, [Av98].

**Proposition 3.1.** *For each formal multigraded power series with integer coefficients*

$$P(y, t) = 1 + \sum_{i=1}^{\infty} \left( \sum_{\mathbf{j}} a_{i,\mathbf{j}} y^{\mathbf{j}} \right) t^i$$

where for each  $i$ ,  $a_{i,\mathbf{j}} = 0$  for all but finitely many values of  $\mathbf{j}$ , there exist uniquely defined integers  $e_{n,\mathbf{j}} \in \mathbb{Z}$  such that

$$P(y, t) = \frac{\prod_{i=1}^{\infty} \prod_{\mathbf{j}} (1 + y^{\mathbf{j}} t^{2i-1})^{e_{2i-1,\mathbf{j}}}}{\prod_{i=1}^{\infty} \prod_{\mathbf{j}} (1 - y^{\mathbf{j}} t^{2i})^{e_{2i,\mathbf{j}}}}$$

and the product converges in the  $(t)$ -adic topology of the ring  $\mathbb{Z}[y, t]$ .

*Proof.* We set  $P_0(t) = 1$  and assume by induction that  $P(y, t) \equiv P_{n-1}(y, t)$  modulo  $t^{n+1}$ . If  $P(y, t) - P_{n-1}(y, t) \equiv \sum e_{n,\mathbf{j}} y^{\mathbf{j}} t^n \pmod{t^{n+1}}$ , then we set  $P_n(y, t) = P_{n-1}(y, t) \prod_{\mathbf{j}} (1 + y^{\mathbf{j}} t^n)^{e_{n,\mathbf{j}}}$  if  $n$  is odd and  $P_n(y, t) = P_{n-1}(y, t) / \prod_{\mathbf{j}} (1 - y^{\mathbf{j}} t^n)^{e_{n,\mathbf{j}}}$  if  $n$  is even. Then it is clear that  $P(t) \equiv P_n(t) \pmod{t^{n+1}}$  and the other assertions follow as well.  $\square$

We define the *multigraded  $(n, \mathbf{j})$  deviation* of  $R$ , denoted by  $\epsilon_{n,\mathbf{j}}$ , to be the exponent  $e_{n,\mathbf{j}}$  in the product decomposition of the Poincaré series of  $R$ . As in Theorem 7.1.3, [Av98] it follows that

**Proposition 3.2.** *Let  $I$  be a monomial ideal of  $S$ ,  $R = S/I$  and  $R\langle X \rangle$  be the multigraded acyclic closure of  $\mathbb{k}$  over  $R$ . Then*

$$P_R(y, t) = \frac{\prod_{i=1}^{\infty} \prod_{\mathbf{j}} (1 + y^{\mathbf{j}} t^{2i-1})^{\text{card}(X_{2i-1,\mathbf{j}})}}{\prod_{i=1}^{\infty} \prod_{\mathbf{j}} (1 - y^{\mathbf{j}} t^{2i})^{\text{card}(X_{2i,\mathbf{j}})}}$$

and  $\text{card } X_{n,\mathbf{j}} = \epsilon_{n,\mathbf{j}}(R)$ .

Since  $R\langle X \rangle$  is multigraded, it follows that the homotopy Lie algebra of  $R$ ,  $\pi^\bullet(R) = H\text{Der}_R^\gamma(R\langle X \rangle, R\langle X \rangle)$  has a multigraded structure. Moreover  $\pi^n(R) \cong \text{Hom}_{\mathbb{k}}(\mathbb{k}X_n, \mathbb{k})$ , (Theorem 10.2.1, [Av98]), and  $\pi_{\mathbf{j}}^n(R) \cong \text{Hom}_{\mathbb{k}}(\mathbb{k}X_{n,-\mathbf{j}}, \mathbb{k})$  while  $\text{rank}_{\mathbb{k}} \pi_{\mathbf{j}}^n(R) = \epsilon_{n,-\mathbf{j}}$ .

**Remark 3.3.** We let  $K_\bullet$  denote the Koszul complex on  $x_1, \dots, x_n$  over  $S$  and  $T_\bullet$  the Taylor complex on the minimal generators of  $I$  over  $S$ . The complexes  $K_\bullet, T_\bullet$  have the structure of a DG  $\Gamma$  algebra by Lemma 9, [Av02]; they are also multigraded and  $H_\bullet(T \otimes_S K)_{\mathbf{j}} \cong H_\bullet(T \otimes_S \mathbb{k})_{\mathbf{j}} \cong H_\bullet(R \otimes_S K)_{\mathbf{j}}$ , where  $\mathbf{j} \in L_I$ . In Lemma 11 of [Av02], it is shown that  $\pi^{\geq 2}(R) \cong \pi^\bullet(R \otimes_S K)$ . We remark that  $R \otimes_S K$  is a multigraded DG  $\Gamma$  algebra, and one can choose a multigraded DG  $\Gamma$

algebra  $U$  so that we have the factorization  $R \otimes_S K \longrightarrow U \longrightarrow \mathbb{k}$  with the properties of Lemma 11 and Remark 10 of [Av02]:  $\pi(R \otimes_S K)$  is the graded  $\mathbb{k}$  dual of the residue of  $H_\bullet(U \otimes_{R \otimes_S K} \mathbb{k})$  modulo multigraded relations and is multigraded. The homomorphisms of Lemma 11, [Av02] are multigraded and  $\pi_j^{l+2}(R) \cong \pi_j^l(R \otimes_S K)$ . Finally we have  $\pi_j^l(T \otimes_S K) \cong \pi_j^l(T \otimes_S \mathbb{k}) \cong \pi_j^l(R \otimes_S K)$ .

Next we will make use of the LCM lattice  $L_I$ , [GaPeWe99] and the GCD graph  $G_I$ , [Av02]. We recall that  $L_I$  and  $G_I$  have vertices the least common multiples of the monomial generators of  $I$  while the edges of  $G_I$  join least common multiples that are relatively prime. In [Av02] it is shown that if  $I$  and  $I'$  are two monomial ideals of  $S$  and  $S'$  respectively with an isomorphism of lattices  $\lambda : L_I \longrightarrow L_{I'}$  which induces an isomorphism of the GCD graphs  $\lambda : G_I \longrightarrow G_{I'}$  then the homotopy Lie algebras of  $R = S/I$  and  $R' = S'/I'$  are isomorphic and the Poincaré series of  $R$  and  $R'$  have the same denominator. We will show that the multigraded version of this result is actually true. We consider the multigraded acyclic closures  $R < X >$  and  $R' < X' >$  of  $\mathbb{k}$  over  $R$  and  $R'$ .

**Remark 3.4.** Let  $I \subset S = \mathbb{k}[x]$  and  $I' \subset S' = \mathbb{k}[x']$  be two monomial ideals,  $K, K'$  the Koszul complexes on the  $x_i$  and  $x'_i$  respectively and  $\lambda : L_I \longrightarrow L_{I'}$  an isomorphism of lattices. If  $\lambda$  induces an isomorphism of the GCD graphs  $\lambda : G_I \longrightarrow G_{I'}$  then  $H(K \otimes R)_j \cong H(K' \otimes R')_{\lambda(j)}$  through the isomorphism that sends the algebra generators  $X_A$  of  $H(K \otimes R) = \mathbb{k}[X_A]/L$  to the generators  $X'_{\lambda(A)}$  of  $H(K' \otimes R') = \mathbb{k}[X'_{\lambda(A)}]/L'$ , (here our notation is as in the remarks preceeding Proposition 2.2). Indeed, let  $\lambda$  be as above. Since  $\lambda$  is an isomorphism of the LCM lattices the minimal resolution of  $I$  gives rise to a minimal resolution of  $I'$  and the generators  $X_A$  of  $H(K \otimes R)$  correspond to generators  $X'_{\lambda(A)}$  of  $H(K' \otimes R')$ . Since  $\lambda$  is an isomorphism of the GCD graphs the relations among the  $X_A$  correspond to relations among the  $X'_{\lambda(A)}$ . Conversely if the Koszul homology algebras are isomorphic as above then there is an isomorphism of multigraded vector spaces between the minimal resolutions of the two ideals and a map  $\lambda$  which is an isomorphism of lattices and GCD graphs for the multidegrees that appear in the resolutions.

**Proposition 3.5.** *Let  $I'$  be a monomial ideal of  $S' = \mathbb{k}[x']$  such that  $\lambda : L_I \longrightarrow L_{I'}$  is an isomorphism of lattices that induces an isomorphism of the GCD graphs  $\lambda : G_I \longrightarrow G_{I'}$ . Then there is a map  $\hat{\lambda}$  from the set of the multidegrees of the variables  $X_n$  to the set of the multidegrees of the variables  $X'_n$  such that  $\text{card } X_{n,j} = \text{card } X'_{n,\hat{\lambda}(j)}$  and  $\epsilon_{n,j} = \epsilon'_{n,\hat{\lambda}(j)}$ .*

*Proof.* We define  $\hat{\lambda}$  by induction on  $n$ . Let  $T'$  be the Taylor complex of  $I'$ . Then there is an isomorphism of vector spaces  $(T \otimes_S \mathbb{k})_j \cong (T' \otimes_{S'} \mathbb{k})_{\lambda(j)}$ , for  $j \in L_I$ . Suppose that  $\hat{\lambda}$  is an extension of  $\lambda$  and  $R[X_{\leq n,j}] \otimes \mathbb{k} \cong R'[X'_{\leq n,\hat{\lambda}(j)}] \otimes \mathbb{k}$ . Extend  $\hat{\lambda}$  to the multidegrees of the multigraded generators of  $H_n(R[X_{\leq n}])$  and  $H_n(R'[X'_{\leq n}])$ . It follows that there is an isomorphism of homotopy Lie algebras  $\pi_{-\hat{\lambda}(j)}^\bullet(T \otimes_S \mathbb{k}) \cong \pi_{-\hat{\lambda}(j)}^\bullet(T' \otimes_{S'} \mathbb{k})$ , and the proposition now follows.  $\square$

**Remark 3.6.** Let  $I$  be a monomial ideal and  $I_{pol}$  be the square free monomial ideal in  $A = \mathbb{k}[z]$  that corresponds to  $I$ , [Fr82]. The map  $\lambda$  that sends a monomial of  $I$  to a squarefree monomial in  $I_{pol}$  is an isomorphism of lattices and of GCD graphs. Moreover  $Q_R(y, t)$  can be obtained from  $Q_A(z, t)$  by applying  $\lambda^{-1}$  to the monomial coefficients of  $t$ , [Ba82].

Theorem 3.7 now completes the multigraded version of Theorem 1, [Av02].

**Theorem 3.7.** *Let  $S = \mathbb{k}[x]$  and  $S' = \mathbb{k}[x']$  be polynomial rings over a field  $\mathbb{k}$ , and  $I \subset S$ ,  $I' \subset S'$  be ideals generated by monomials of degree at least 2. Let  $\lambda : L_I \rightarrow L_{I'}$  and the induced  $\lambda : G_I \rightarrow G_{I'}$  be isomorphisms. Suppose that  $Q_R(y, t) = 1 + \sum (\sum c_j y^j) t^i$  then  $Q_{R'}(y', t) = 1 + \sum (\sum c_j y^{\lambda(j)}) t^i$ .*

*Proof.* By Proposition 2.1 the multidegrees  $\mathbf{j}$  in  $Q_R(y, t)$  are in  $L_I$ , so  $\lambda(\mathbf{j})$  makes sense. By the above remark it is enough to examine the case where  $I$  and  $I'$  are both squarefree. The key point is that since  $I$  and  $I'$  are squarefree,  $\lambda$  is additive:  $\lambda(\mathbf{j} + \mathbf{i}) = \lambda(\mathbf{j}) + \lambda(\mathbf{i})$ . It follows that its extension  $\hat{\lambda}$  is also additive. Since

$$Q_R(y, t) = \frac{\prod_{i=1}^n (1 + ty_i)}{P_R(y, t)} = \frac{\prod_{i=1}^{\infty} \prod_{\mathbf{j}} (1 - y^{\mathbf{j}} t^{2i})^{\text{rank } \pi_{-\mathbf{j}}^{2i}}}{\prod_{i=2}^{\infty} \prod_{\mathbf{j}} (1 + y^{\mathbf{j}} t^{2i-1})^{\text{rank } \pi_{-\mathbf{j}}^{2i-1}}}$$

the linear coefficient  $c_j$  of the monomial  $y^j t^n$  of  $Q_R(y, t)$ , depends on the deviations  $\epsilon_{i, \mathbf{f}}$  when  $i \leq n$ . The corresponding combination of the deviations  $\epsilon'_{i, \hat{\lambda}(\mathbf{f})}$  gives a term of  $Q_{R'}(y', t)$  of  $t$ -degree  $n$ . The theorem now follows.  $\square$

The condition that  $\lambda$  is an isomorphism of GCD graphs is a necessary condition as the example from [Ga00] shows. The isomorphism of LCM lattices for  $I = (x_1^2, x_2^2 x_3)$  and  $I' = (x_1 x_2^2, x_1 x_3^2)$  is not an isomorphism of GCD graphs and

$$Q_{S/I}(y, t) = 1 - t^2(y_1^2 - y_2^2 y_3) + t^4(y_1^2 y_2^2 y_3)$$

while

$$Q_{S/I'}(y, t) = 1 - t^2(y_1 y_2^2 + y_1 y_3^2) - t^3(y_1 y_2^2 y_3^2).$$

## REFERENCES

- [Av98] L. AVRAMOV *Infinite Free Resolutions*, in: Six Lectures in Commutative Algebra, Bellaterra, 1996, Progress in Math. **166**, Birkhäuser, Boston, (1998), 1-118
- [Av02] L. AVRAMOV *Homotopy Lie Algebras and Poincaré series of Algebras with Monomial Relations*, Homology Homotopy Appl. **4**(2), (2002), 17-27
- [Ba82] J. BACKELIN *Les anneaux locaux à relations monomiales ont des séries de Poincaré-Betti rationnelles*. C.R. Acad. Sc. Paris, Série I, **295** (1982), 607-610.
- [BaPeSt98] D. BAYER, I. PEEVA, AND B. STURMFELS *Monomial resolutions*, Mathematical Research Letters, **5** 1998, 31-46
- [ChRe95] H. CHARALAMBOUS AND A. REEVES *Poincaré Series and Resolutions of the Residue Field over Monomial Rings*, Com. Algebra, **23**(6), 1995, 2389-2399
- [Ei97] D. EISENBUD *Commutative Algebra with a View Toward Algebraic Geometry*, Springer Verlag, 1994
- [Fr79] R. FRÖBERG *Some complex constructions with applications to Poincaré series*, Séminaire d'algèbre P. Dubreil 1977/78, Lecture Notes in Mathematics, Springer-Verlag, **740**, (1979), 272-284.
- [Fr82] R. FRÖBERG *A study of graded extremal rings and of monomial rings*, Math. Scand. bf 51, (1982), 22-34
- [Ga00] V. GASHAROV *Infinite Monomial Resolutions*, Comm. Algebra, **28**(11), (2000), 5455-5460
- [GaPeWe99] V. GASHAROV, I. PEEVA, AND V. WELKER *The LCM-Lattice in Monomial Resolutions*, Math. Research Letters, **6**, (1999), 521-532
- [GuLe69] T.H. GULLIKSEN AND G. LEVIN *Homology of local rings*, Queen's Paper no. 20, Queen's University, Kingston (1969).
- [Iy97] S. IYENGAR *Free resolutions and change of rings*, J. Algebra, **190**, (1990), pp 195-213.

- [Le86] J. LESCOT *Séries de Poincaré des modules multi-gradués sur les anneaux monomiaux* Algebraic topology-rational homotopy, Lect. Notes in Math., Springer, **1318** (1988), 155-161
- [MiStYa00] E. MILLER, B. STURMFELS, AND K. YANAGAWA *Generic and Cogeneric monomial ideals*, Symbolic Computation in Algebra, Analysis, and Geometry (Berkeley, CA, 1998), J. Symbolic Comput., **29** (2000), 691–708.
- [Ta60] D. TAYLOR *Ideals generated by monomials in an R-sequence*, Thesis, University of Chicago, 1960.

DEPARTMENT OF MATHEMATICS, UNIVERSITY AT ALBANY, SUNY, ALBANY, NY 12222  
*E-mail address:* `hara@math.albany.edu`